



Black-crowned Night-Heron

Nycticorax nycticorax



Joe Kosack/PGC Photo

CURRENT STATUS: In Pennsylvania, the black-crowned night-heron is listed as state endangered and protected under the Game and Wildlife Code. Nationally, they are not listed as an endangered/threatened species. All migratory birds are protected under the federal Migratory Bird Treaty Act of 1918.

POPULATION TREND: The black-crowned night-heron (*Nycticorax nycticorax*) was once considered a fairly common nesting heron in Pennsylvania. In fact, B. H. Warren (1888) claimed it was the most common nesting heron in the state next only to the green heron (*Butorides virescens*). Black-crowned night-herons have always been more common in the southeastern counties than in other parts of the state, mainly along the Susquehanna and Delaware River corridors, but the species is now difficult to find anywhere else in the state. The number and size of colonies have declined in recent years, especially in the northern and western counties (Luzerne and Crawford, for example) where the species no longer nests.

The Pennsylvania Game Commission and agency partners count active nests at known nest sites every year to monitor population trends. In 2009, black-crowned night-herons were found nesting in five colonies statewide: Wade Island, Dauphin County; Ephrata, Lancaster County; Kiwanis Lake, York County; Kinzer, Lancaster County; and Wyomissing Hills, Berks County. Wade Island and Kiwanis Lake are Pennsylvania Audubon Important Bird Areas. The Wade Island colony has been surveyed annually since 1985. It reached its peak in 1990 with 345 nesting pairs and has declined since that year. The 2007, 2008 and 2009 surveys counted 62, 98 and 116 nests respectively. The colony currently is above the 10-year (1999-2008) average of 99 nests. The Kiwanis Lake colony has been surveyed for the past decade and there are indications competition for nest trees may be limiting the colony's growth. Also, many of these pairs are dispersing to adjacent neighborhoods or lakes as nest trees age. There were 117 nests in 2007; 100 nests in 2008; and 122 in 2009. Combined, the Kiwanis Lake and Wade Island colonies comprise almost 90 percent of the state's population of black-crowned night-herons. The Ephrata colony has existed since 2006 and is stable to increasing with several dozen pairs. The other colonies are small, averaging four or five nests in each.



Joe Kosack/PGC Photo

IDENTIFYING CHARACTERISTICS: This medium-sized heron is very stocky and short-necked in appearance. The black-crowned night-heron is slightly larger than an American crow (1.6 - 2.2 pounds versus 1 - 1.4 pounds; Davis 1993), but less than half the size of the more familiar great blue heron (4.6 - 5.5 pounds; Butler 1992). The adults are distinctly marked with a black cap and upper back, contrasting with gray wings, rump and tail and white to light-

gray breast. It is known by some as “the squawk” for its distinctive croaking call given at dusk as it flies to its evening feeding place. Juveniles have a streaky brown plumage with large white spots on the wing coverts. By contrast, the yellow-crowned night-heron is a smaller (1.3 – 1.8 pounds; Watts 1995), more delicate-looking heron with longer neck and legs.




Joe Kosack/PGC Photo

BIOLOGY-NATURAL HISTORY: Night-herons are very social; they nest in colonies and usually forage together. As their name suggests, night-herons are chiefly nocturnal and crepuscular in habit (that is, they are active at night or at dawn and dusk). They build simple stick nests in trees near where they hunt for food in shallow waters, opportunistically searching for a wide variety of prey items. If free of predators or human interference, they can maintain a colony for decades. They may not use a nest every year, but they may return to old nests after an extended absence. Night-herons are opportunistic predators, foraging for foods such as small fishes, crustaceans such as crayfish, insect larvae, leeches, mussels, mice, small birds, reptiles and amphibians, various plant materials, and carrion. Like most herons, black-crowned night-herons will take advantage of fish and crayfish hatcheries for food. Its high level on the food

chain, colonial nesting, and wide distribution throughout its range make this species a good indicator of environmental quality and vulnerable to environmental contaminants.

Nesting can occur early in Pennsylvania (March-April) and continue through June or July, particularly if bad weather caused an early nest to fail. Black-crowned night-heron clutches throughout North America average 3-5 eggs (range 1-7) and are greenish in color (Davis 1993). Adults share the 24- to 26-day incubation period, sometimes fighting over trading incubation duties. They also share responsibilities of raising the chicks to fledging, six to seven weeks after hatching. Because the parents do not recognize their own chicks for a couple of months, juveniles can abandon the nest and continue to beg for food from any adult black-crowned night-heron in the colony. Juveniles will ultimately leave nesting areas with adults to travel to foraging areas. This dispersal following the breeding season allows for early morning or evening viewing opportunities along streams, lakes or ponds. By mid-September, birds travel to their southern wintering areas along the eastern and Gulf coasts of the United States and into Central America. Wintering birds have been observed at John Heinz National Wildlife Refuge (Philadelphia, PA) in small numbers (see McWilliams and Brauning 2000 for other locales).

PREFERRED HABITAT: The black-crowned night-heron is fairly widespread in distribution. Its nesting range reaches north into the Great Lakes, St. Lawrence valley and the coast of New England. Its habitat is quite varied, but always includes water. They can be found along coasts, along streams, in swamps, and in some human-altered locations such as canals and wet agricultural fields. Night-herons nest in trees, including native riparian species (sycamores, maples, ashes, birches) and exotic conifers. In Pennsylvania, it is found primarily around rivers and creeks, often nesting on forested islands or along wooded streams. They can be fairly tolerant of human habitations and, if protected, will nest close to houses and in towns and cities. Proximity to humans may afford them some protection from predators. Dispersal following the breeding season can be to the north before they travel south to overwinter in warmer climates. So, many night-herons are observed in northern Pennsylvania after the nesting season is finished.



REASONS FOR BEING ENDANGERED: The decline in the number and size of Pennsylvania colonies is the main reason this species has been listed as "endangered" in the state. The remaining colonies are few and vulnerable to either human-caused or natural destruction. The state's largest colony on Wade Island in the Susquehanna River near Harrisburg is threatened by erosion and competition for nest sites with great egrets (*Ardea alba*) and a recent colonizer, the double-crested cormorant (*Phalacrocorax auritus*). Population declines are not well-understood. Wetland losses in southern Pennsylvania may be an important factor in its decline. In the past, plumage hunters took a significant toll of nesting herons. Many waterbirds were hunted for their feathers because it was popular for women to adorn their hats with feathers and entire birds. Little is known about the impact of feral cats and dogs or nest predators such as raccoons and crows. Shooting birds at fish hatcheries, legal and illegal, is another possible factor in population declines. Little is known about the foraging range and preferences of the state's heron colonies so those vulnerabilities are difficult to assess.

MANAGEMENT PROGRAMS: Nesting colonies are protected through the Pennsylvania Natural Heritage Program and the Environmental Review process. Colonies are monitored through the PGC's colonial waterbird program. The largest colonies are part of the Sheets Island Archipelago and Kiwanis Lake Important Bird Areas. More studies of the ecology and behavior of this species in Pennsylvania are needed to better understand its conservation needs.

Sources:

Davis, W. E. , Jr. 1993. Black-crowned night-heron (*Nycticorax nycticorax*). In The Birds of North America, No. 74 (A. Poole and F. Gill, Eds.). Philadelphia: The Academy of Natural Sciences; Washington D. C.: The American Ornithologists' Union.



McWilliams, G. M. and D. W. Brauning. 2000. The Birds of Pennsylvania. Cornell University Press, Ithaca, NY.

Warren, B. H. 1888. Report on the Birds of Pennsylvania. State Board of Agriculture, Harrisburg, PA.

Suggested further reading:

Crossley, G. J. (Compiler). 1999. A Guide to Critical Bird Habitat in Pennsylvania: Pennsylvania Important Bird Areas Program. Pennsylvania Audubon Society, Harrisburg, PA.

Hancock, J. and H. Elliot. 1978. The Herons of the World. Harper & Row, Publishers, New York, NY. p. 124-128.

NatureServe. 2009. [NatureServe Explorer](#): An online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Search for "black-crowned night-heron."

Pennsylvania Audubon. 2005. [Important Bird Area Conservation Plans](#). Pennsylvania Audubon Society, Harrisburg, PA.

Schutsky, R. M. 1992. The Black-crowned Night-Heron. In Atlas of the Breeding Birds of Pennsylvania (D. W. Brauning, Ed.). University of Pittsburgh Press, Pittsburgh, PA. pp. 60-61.

By Cathy Haffner and Doug Gross
Pennsylvania Game Commission
Connecting you with wildlife!

10/7/09