Topic Area TU: Tobacco Use

Objective TU-6: **Percent Smoke Cessation During First Trimester of Pregnancy* - 2020 Target: 30.0**

Percent Smoke Cessation During First Trimester of Pregnancy* by Race and Hispanic Origin, Pennsylvania Residents, 2007-2011



Smoke Cessation During First Trimester of Pregnancy*, Percent and Number for Selected Demographics, Pennsylvania Residents, 2007-2011

	PENNSYLVANIA									
	2007		2008		2009		2010		2011	
	%	No.	%	No.	%	No.	%	No.	%	No.
Smoke cessation during first trimester of pregnancy* mothers 18-49	25.0	8,035	25.2	7,882	25.6	7,733	26.2	7,597	26.3	7,378
White mothers 18-49	25.6	6,737	25.6	6,565	26.0	6,388	26.6	6,271	26.3	6,024
Black mothers 18-49	18.8	713	19.6	719	19.9	724	20.8	728	22.3	774
Hispanic mothers 18-49	30.2	558	31.1	560	28.8	534	32.9	598	33.1	577
Asian/Pacific Islander mothers 18-49	34.4	66	36.5	73	48.8	100	47.9	80	47.5	85

*Mothers counted as quitting smoking were those who smoked during the three months prior to the pregnancy and did not smoke during the first, second, and third trimesters.

NOTE: Hispanics can be of any race.
The above statistics and additional data for Healthy People 2020 objectives can also be found at www.health.state.pa.us/stats (select Healthy People).
DSU = Data statistically unreliable (fewer than 10 events) NE = No events